

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**




FIG. 1

2/10




FIG. 2




FIG. 3





FIG. 4

Original gel spot #	Protein Identity	Function	Venlafaxine (Fold change)	Fluoxetine (Fold change)	Accession#	MOWSE score	Protein Area Coverage	Mr/pI	Species
87	IGF-1 A precursor	GH is an important regulator of IGF-1 expression. Secreted/Growth-promoting activity.	2.9	2.5	P08025	7.07E+01	28%	17079/9.5	Rat

FIG. 5


6/10

Differential Expression of IGFBP2 mRNA: Amygdala Long-Term



Treatment (14 Days)

Differential Expression of IGFBP2 mRNA: Amygdala Short-Term



Treatment (3 Days)

FIG. 6






FIG. 7

8/10



NBI-31772

(Liu et al, Neurocrine Biosciences, JBC 276:32419, 2001)

FIG. 8

FIG. 9




FIG. 10